

The method for transforming immature wheat embryos was developed and optimized by Becker and Lörz (D. Becker and H. Lörz, Plant Tissue Culture Manual (1996), B12: 1 to 20).

- 5 In the experiments described hereinbelow, the procedure developed by Becker and Lörz (loc. cit.) was adhered to.

For the transformation, ears with caryopses of developmental stage 12 to 14 days after anthesis were harvested and surface-sterilized. The isolated
10 scutella were plated onto induction medium #30 with the embryo axis orientated towards the medium.

After preculture for 2 to 4 days (26°C, in darkness), the explants are transferred to medium #39 for the osmotic preculture (2 to 4 h, 26°C, in the
15 dark).

For the biolistic transformation, approx. 29 µg of gold particles onto which a few µg of the target DNA had previously been precipitated were employed per shot. Since the experiments carried out are cotransformations, the
20 target DNA added to the precipitation batch is composed of the target gene and a resistance marker gene (bar gene) in the ratio 1:1.

4. DIG labeling of DNA fragments

25 DNA fragments employed as screening probes were labeled via a specific PCR with the incorporation of DIG-labeled dUTP (Boehringer Mannheim, Germany).

Media solutions used in the examples:

30

20 × SSC	175.3 g NaCl
	88.2 g sodium citrate
	twice-distilled H ₂ O to 1000 ml
	10 N NaOH to pH 7.0

35

Plasmid pTaSSI 8/1 was deposited at the DSMZ in Braunschweig, Federal Republic of Germany, as specified in the Budapest Treaty under the

009011-12842960